

III. Topology and Hodge theory

- ▶ These two topics are closely intertwined and constitute a major aspect of complex algebraic geometry, beginning in the later part of the 19th century (Picard, Poincaré, ...) into the 1st half of the 20th century (Lefschetz, Hodge, ...) and continuing through today
- ▶ In fact questions about integrals on algebraic surfaces (which are real 4-manifolds) were instrumental in the beginnings of topology — one knew (Darboux, Picard, Poincaré, E. Cartan, ...) what differential forms

$$\varphi = a \, dx + b \, dy + c \, dz$$

$$\psi = A \, dx \wedge dy + B \, dx \wedge dz + C \, dy \wedge dz$$

$$\eta = D \, dx \wedge dy \wedge dz$$

were, and Stokes' theorem

$$\int_U d\omega = \int_{\partial U} \omega$$

shows then when $d\omega = 0$ that $\int_{\Gamma} \omega$ was not only invariant under deformation (homotopy) of Γ but also under homology.¹ This led to the notion of *periods*

$$\int_{\Gamma} \omega, \quad d\omega = 0 \text{ and } \Gamma \in H_p(X, \mathbb{Z}).$$

¹The exterior derivative d is uniquely determined (i) $df = f_x dx + f_y dy + f_z dz$ for a function f , (ii) $d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^{\deg \alpha} \alpha \wedge d\beta$ and (iii) $dx \wedge dy = -dy \wedge dx$ etc.

In the complex case when X has local holomorphic coordinates $z = (z_1, \dots, z_n)$

$$\omega = \sum_{I,J} f_{I\bar{J}} dz^I \wedge d\bar{z}^J$$

where $I = (i_1, \dots, i_p)$, $dz^I = dz^{i_1} \wedge \dots \wedge dz^{i_p}$ etc. and as we saw for algebraic curves the periods reflect the complex structure — this is the start of Hodge theory.

Outline for the remainder of this lecture

- ▶ Introductory discussion of what an algebraic variety is
- ▶ Statements of the Lefschetz theorems
- ▶ How they arose historically from the study of algebraic functions of two variables (Picard-Lefschetz or PL theory)
- ▶ Origin of the Hodge conjecture (HC)



- ▶ *Complex projective space* $\mathbb{P}^N$
 - ▶ lines through origin in $\mathbb{C}^{N+1}$
 - ▶ $\mathbb{P}^N = \mathbb{C}^N \cup \mathbb{P}^{N-1}$ ($\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$)
 - ▶ homogeneous coordinates $[z] = [z_0, \dots, z_N]$
- ▶ $\mathbb{P}^1$ = Riemann sphere
- ▶ $\mathbb{P}^2 = \mathbb{C}^2 \cup \{\text{lines through the origin}\}$ where $[z] \leftrightarrow$ line with slope z_2/z_1
- ▶ $\mathbb{P}^N$ = compact complex manifold

Proof $\mathcal{U}_i = \{[z] : z_i \neq 0\} \ni [z]$

$\downarrow$

$\downarrow$

$$\mathbb{C}^N \ni (z_0/z_i, \dots \overset{i}{\wedge} \dots, z_N/z_i)$$

- ▶ Algebraic variety $X \subset \mathbb{P}^N$ given by $F_1(z) = \dots F_m(z) = 0$ where $F_\alpha(z)$ = homogeneous polynomial.
- ▶ Note that $\dim_{\mathbb{R}} X = 2 \dim_{\mathbb{C}} X$ and X is oriented.

Example


C defined by $f(x, y) = 0$ in $\mathbb{C}^2$. Set

$$x = z_1/z_0, \quad y = z_2/z_0$$

and clear denominators to get



$$\overline{C} = \{F(z) = 0\} \subset \mathbb{P}^2$$

$$\text{where } \overline{C} = \left\{ \begin{array}{l} \text{our old} \\ C \subset \mathbb{C}^2 \end{array} \right\} \cup \left\{ \begin{array}{l} \text{points} \\ \text{at } \infty \end{array} \right\}.$$




- ▶ suppose X^n = smooth algebraic variety and $Y = \mathbb{P}^{N-1} \cap Y$ is a general hyperplane section

hyperplane section



quadric surface;
real picture



Note: Equation of the quadric in $\mathbb{C}^3$ is $x^2 + y^2 = z^2 + 1$; equation in $\mathbb{P}^3$ is $z_1^2 + z_2^2 = z_3^2 + z_0^2$; over $\mathbb{C}$ this is equivalent to $z'_1 z'_2 = z'_3 z'_0$ where $z'_1 = z_1 + iz_2$, $z'_2 = z_1 - iz_2$ etc.

Lefschetz theorem I

- ▶ $b_{2p+1}(X) \equiv 0 \pmod{2}$ (odd Betti numbers are even)
- ▶ $b_{2p}(X) \geq 1$ (even Betti numbers are positive).

In the second, if $\dim_{\mathbb{C}} X = n$ and $H \in H_{2n-2}(Y, \mathbb{Z})$ is the class of the cycle given by Y then (non-trivially)

$$\underbrace{H \cap \cdots \cap H}_{n-p} \neq 0 \text{ in } H_{2p}(X, \mathbb{Z})$$

Lefschetz theorem II


$$H_p(Y, \mathbb{Z}) \rightarrow H_p(X, \mathbb{Z}) \text{ is } \begin{cases} \text{isomorphism for} \\ p \leq n-2 \end{cases}$$

onto for $p = n-1$

Corollary

Y is connected if $\dim_{\mathbb{C}} X \geq 2$

Exercise: $f(x, y)$ = irreducible polynomial and $\{f(x, y) = 0\} = C \subset \mathbb{C}^2$. Show that C is connected.




Geometric idea to study topology of an algebraic variety (idea is one of the most basic in algebraic geometry) — use induction by dimension.


Example

For $y^2 = p(x)$ where $p(x) = \prod_{i=1}^{2g+2} (x - a_i)$

- ▶ first take out the two points over $x = \infty$
- ▶ next use the picture of the complex x -plane



- ▶ retract the slit x -plane and the part of C lying over it onto the part lying over the segments

▶  $= \left\{ \begin{array}{l} \text{1-dimensional} \\ \text{complex} \end{array} \right\}$

- ▶ on  as we turn around the branch point the two points interchange (local *monodromy* T_i around a_i)

- ▶ $\prod_i T_i = \text{Id}$

- ▶ $\Delta_i = \begin{array}{c} + \\ \bullet \\ \diagup \quad \diagdown \\ - \bullet \end{array}$ lying over $\begin{array}{c} \bullet \quad \bullet \\ 0 \quad a_i \end{array}$

- ▶ C retracts onto the real 1-dimensional complex given by attaching the $2g + 2$ 1-cells Δ_i to the two points $\pm$ lying over 0.
- ▶ Δ_i generate the relative homology group
 $H_1(C, \{+, -\}; \mathbb{Z})$
 $\rightsquigarrow H_1(\overline{C}, \mathbb{Z}) \cong \mathbb{Z}^{2g}$

This case is too simple to suggest the general pattern. The next dimension up is due to Picard (1880–2000)

Example


X is the algebraic surface

$$z^2 = f(x, y)$$

where $C = \{f(x, y) = 0\}$ is a non-singular plane curve. For a general y we let

$$X_y = \text{curve } z^2 = f(x, y), \quad y \text{ fixed}$$


The picture is



X_y is the algebraic curve of the type we have been considering; it is 2:1 covering of the line $y = \text{constant}$ branched at the points of $C \cap \{y = \text{constant}\}$

- ▶ smooth for general y
- ▶ singular when the line $y = \text{constant}$ becomes tangent to C

- ▶ the picture of X_y is




where the branch points and slits will vary with y

- ▶ at a point of tangency two branch points come together and interchange.

► $\delta \rightarrow \delta$

► $\gamma \rightarrow ?$



Picard-Lefschetz formula

(PL)

$$\gamma \rightarrow \gamma + \delta$$


How to show PL? The original argument was analytic and in outline went as follows:

- ▶ locally analytically change coordinates so that the picture is a neighborhood of the origin of the curves

$$C_t = \{u^2 + v^2 = t\}$$

in $\mathbb{C}^3$ with coordinates (u, v, t)

- ▶ the local picture is



- ▶ set $t = \sigma^2$ and consider the integrals

$$I_t(\delta) = \int_{\delta} \frac{du}{\sqrt{t - u^2}} = \int_{\delta} \frac{du}{\sqrt{\sigma^2 - u^2}} = \int_{\delta} \frac{du}{\sigma \sqrt{1 - (u/\sigma)^2}}$$


$$I_t(\gamma) = \int_{\gamma} \frac{du}{\sqrt{t - u^2}} = \int_{\gamma} \frac{du}{\sqrt{\sigma^2 - u^2}} = \int_{\gamma} \frac{du}{\sigma \sqrt{1 - (u/\sigma)^2}}$$

- ▶ the curves C_t are parametrized by

$$z \rightarrow (\sigma \sin z, \sigma \cos z),$$

and a calculation gives

$$\begin{cases} I_t(\delta) = 2\pi \\ I_t(\gamma) = i \log t \end{cases}$$



Conclusion


$$\begin{cases} l_{e^{2\pi i t}}(\delta) = l_t(\delta) \\ l_{e^{2\pi i t}}(\gamma) = l_t(\gamma) + l_t(\delta) \end{cases}$$

$$\implies T(\gamma) = \gamma + \delta.$$




Topological pictures


global



local




$$\delta \rightarrow 0$$




- ▶ few pictures worth 1,000 (10,000?) words
- ▶ heuristic analytic reasoning suggests what the answer should be — then know what to prove.

- ▶ $X^* = X \setminus X_\infty$
- ▶ topological picture of X^*



- ▶ along $\overline{y_0 y_i}$ we have the locus of the vanishing cycle



$\implies$ ▶ X^* obtained from X_0 by attaching 2-cells Δ_i

- ▶ In general

X^* obtained from X_0 by attaching
 $n = \frac{1}{2} (\dim_{\mathbb{R}} X)$ cells

$\implies$ Lefschetz theorems I, II

- ▶ *Single and double integrals*

Returning to X given by

$$z^2 = f(x, y)$$

there are single integrals (1-forms)

$$\psi = \frac{p(x, y) dx}{z} + \frac{q(x, y) dy}{z}$$

and double integrals (2-forms)

$$\varphi = \frac{r(x, y) dx \wedge dy}{z}$$

The story of the ψ 's is very interesting but we will only have time to make a few observations. For one such we note that

$$\blacktriangleright \int \psi < \infty \implies d\psi = 0.$$

Proof:

$$\begin{aligned} d\psi &= d\left(\frac{p(x,y)}{z}\right) \wedge dx + d\left(\frac{q(x,y)}{z}\right) \wedge dy \\ &= \frac{r(x,y) dx \wedge dy}{z} \end{aligned}$$

$$\begin{aligned} \implies \frac{1}{4}(d\psi \wedge \overline{d\psi}) &= \left|\frac{r(x,y)}{z}\right|^2 \left(\frac{i}{2}\right) dx \wedge d\bar{x} \wedge \left(\frac{i}{2}\right) dy \wedge d\bar{y} \\ &= \text{volume form on } X \end{aligned}$$

$$0 < \int_X d\psi \wedge \overline{d\psi} = \int_X d(\psi \wedge \overline{d\psi}) = 0 \implies d\psi = 0.$$

- ▶ The space of single integrals is denoted by $H^{1,0}(X)$ and its dimension $h^{1,0}(X)$ is called the *irregularity* — reason for the name is that in the early days “most” surfaces seemed to be *regular*, i.e., to have $h^{1,0}(X) = 0$.

Example

For $z^2 = f(x, y)$ to be irregular the curve C cannot be smooth, or even have generic singularities, those being where

$$\begin{cases} f_x(x, y) = f_y(x, y) = 0 \\ \det \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix} (x, y) \neq 0 \end{cases}$$

Similarly for a hypersurface

$$F(z_0, z_1, z_2, z_3) = 0$$

in $\mathbb{P}^3$ it is not easy to write down on F where X is irregular.

- Suppose now φ is a regular 2-form; i.e.,

$$\int_{\sigma} \varphi < \infty$$

for any 2-chain σ . We set

$$H^{2,0}(X) = \left\{ \begin{array}{c} \text{space of} \\ \text{regular 2-forms} \end{array} \right\}.$$

The *periods* of ψ are the

$$\int_{\Gamma} \psi, \quad \Gamma \in H_2(X, \mathbb{Z}).$$

Among the Γ 's are the fundamental classes of algebraic curves $C \subset X$; i.e., the images of

$$H_2(C, \mathbb{Z}) \rightarrow H_2(X, \mathbb{Z}).$$

We will discuss these further below.

- By restriction

$$\psi \rightarrow \psi_y = \frac{p(x, y) dx}{z}$$

we will generally have $\psi_y \neq 0$ which gives

$$H^{1,0}(X) \hookrightarrow H^{1,0}(X_y).$$

This suggests that we have

$$H^1(X, \mathbb{C}) \hookrightarrow H^1(X_y, \mathbb{C}),$$


which is true and is what originally suggested the first non-easy case of Lefschetz II — again analysis and topology went hand in hand.

Another example of the use of analysis to suggest topology:

For a vanishing cycle



traced out by $\delta_y \in H_1(X_y, \mathbb{Z})$ along the path from 0 to a_i



we have

$$\int_{\delta_0} \psi = \int_{\delta_{a_i}} \psi = 0, \quad \psi \in H^{1,0}(X).$$

This led to Picard's argument that


$$\ker\{H_1(X_0, \mathbb{Z}) \rightarrow H_1(X, \mathbb{Z})\} = \begin{cases} \text{span of the} \\ \text{space of vanishing cycles.} \end{cases}$$



Returning to the discussion of

- ▶ Among the classes in $H_2(X, \mathbb{Z})$ are those given by the fundamental classes of the algebraic curves C contained in X .

Example:



$$\rightsquigarrow H_2(X, \mathbb{Z}) \cong \mathbb{Z}[L_1] \oplus \mathbb{Z}[L_2]$$

► In general C is a component of

$$\begin{cases} z^2 = f(x, y) \\ g(x, y, z) = 0 \end{cases}$$

(may take $g(x, y, z) = g_0(x, y) + g_1(x, y)z$)

²The lines are $z_0 = z_2 = 0$, $[z_1, z_3] \in \mathbb{P}^1$ arbitrary and $z_1 = z_3 = 0$, $[z_0, z_2]$ arbitrary.

$\Rightarrow$ On X

$$0 = dg = g_x dx + g_y dy + g_z dz$$

which using $dz = \left(\frac{1}{2z}\right) (f_x dx + f_y dy)$ gives a relation

$$a dx + b dy \big|_C = 0$$

$$\Rightarrow \psi \big|_C = 0$$

$$\Rightarrow \int_{[C]} \psi = 0.$$

Conclusion: *The periods of $H^{2,0}(X)$ on the homology classes of algebraic curves are equal to zero.*

- ▶ The converse statement is the famous *Lefschetz (1,1) theorem*.
- ▶ The converse to the analogous statement for arbitrary X is the *Hodge conjecture*.
- ▶ In terms of differential forms of degree 2 on X there are three types:
 - ▶ $\frac{p(x,y) dx \wedge dy}{z} \leftrightarrow H^{2,0}(X)$
 - ▶ conjugates of these $\leftrightarrow \overline{H^{2,0}(X)} = H^{0,2}(X)$
 - ▶ those that have a $dx \wedge d\bar{x}, dx \wedge d\bar{y}, d\bar{x} \wedge dy, dy \wedge d\bar{y}$ which are said to be of type (1,1) and contribute $H^{1,1}(X)$ to $H^2(X, \mathbb{C})$; it is these that are Poincaré dual to the homology classes carried by the algebraic curves in X .

Further topics

- ▶ These involve the *multiplicative structure* on cohomology:
For X of dimension n and $H \in H^2(X)$ the class of a hyperplane section

$$(*) \quad L^k : H^{n-k}(X) \rightarrow H^{n+k}(X).$$

Hard Lefschetz theorem: $(*)$ is an isomorphism

Lefschetz stated the result but his proof was incomplete.
Hodge developed Hodge theory to prove $(*)$.

- ▶ Define operators L, H, Λ on $H^*(X)$ by
 - ▶ L as above
 - ▶ $H = (d - n) \text{Id}$ on $H^d(X)$

Then the commutator

$$[H, L] = 2L.$$

There is a unique $\mathfrak{sl}_2 = \{L, H, \Lambda\}$ with

$$\begin{cases} [L, \Lambda] = H \\ [H, \Lambda] = -2\Lambda. \end{cases}$$

Decomposing $H^*(X)$ into irreducible $\mathfrak{sl}_2$ -modules gives the *Lefschetz decomposition* of cohomology into primitive subspaces — every class is a linear combination of powers of L applied to primitive classes

$$\begin{cases} L^k \cdot \eta \\ \Lambda \eta = 0. \end{cases}$$

- ▶ Any irreducible $\mathfrak{sl}_2$ -module is isomorphic to
 - ▶ $V = \text{span}\{x^n, x^{n-1}y, \dots, xy^{n-1}, y^n\}$
 - ▶ $L = \partial_x, \Lambda = \partial_y$
 - ▶ primitive part is generated by x^n .

Example: $X =$ algebraic surface

$$H^1(X) \xrightarrow{\sim} H^3(X)$$

and

$$H^0(X) \xrightarrow{L} H^2(X) \xrightarrow{L} H^4(X)$$

has

- ▶ $H^2(X)_{\text{prim}} = \ker\{H^2(X) \xrightarrow{L} H^4(X)\}$
- ▶ $H^2(X) = LH^0(X) \oplus H^2(X)_{\text{prim}}$

- ▶ Finally, you may ask: OK, we know a lot about the homology of X — what about its homotopy?

Theorem

The rational homotopy type of X is uniquely determined by $H^(X)$.*

Thus the

- ▶ $\pi_i(X) \otimes \mathbb{Q}$
 - ▶ Massey triple products $/\mathbb{Q}$, etc. are all equal to zero
- ⇒ Very strong homotopy-theoretic conditions that X be topologically a smooth algebraic variety.

Appendix: Monodromy

- ▶ C_0 = smooth algebraic curve over the origin



- ▶ fundamental group $\pi_1 = \pi_1(\mathbb{C} \setminus \{\text{slits}\})$ acts on $H_1(C_0, \mathbb{Z})$
- ▶ action of π_1 is generated by PL transformation

$$T_i : \gamma \rightarrow \gamma + (\gamma, \delta_i) \delta_i$$

- ▶ $\prod T_i = \text{identity}$
- ▶ action of π_1 preserves the intersection form

$$Q : H_1(C_0, \mathbb{Z}) \otimes H_1(C_0, \mathbb{Z}) \rightarrow \mathbb{Z}$$

- ▶ Invariant cycles

$$H_1(C_0, \mathbb{Q})^{\text{inv}} = \text{span}\{\gamma : (\gamma, \delta_i) = 0 \text{ for all } i\}$$

- ▶ Vanishing cycles

$$H_1(C, \mathbb{Q})^{\text{van}} = \text{span}\{\delta_i\}$$

- ▶ If we know that

$$(*) \quad H_1(C_0, \mathbb{Q})^{\text{van}} \cap H_1(C_0, \mathbb{Q})^{\text{inv}} = (0)$$

then

$$Q = \begin{pmatrix} * & * = 0 \\ 0 & * \end{pmatrix}$$

and the monodromy representation is semi-simple

- ▶ Lefschetz stated $(*)$ but his proof was incomplete — in fact

$(*)$ is true, but its proof requires analysis

The analysis was provided by Hodge.

- ▶ It is a general fact proved by Deligne in the geometric case and by Schmid in general that *general monodromy representations are always semi-simple*.

The proofs require Hodge theory and are among the most basic properties of the topology of algebraic varieties.

- ▶ The reason Lefschetz wanted to have the result is that

$$(*) \iff \text{Hard Lefschetz}$$

Lefschetz proof of this assertion was correct.

Hodge structure of weight n

(V, F^p)

$$\left\{ \begin{array}{l} V = \mathbb{Q}\text{-vector space} \quad (\text{frequently have } V_{\mathbb{Z}}) \\ F^p = \text{decreasing filtration on } V_{\mathbb{C}}, \quad 0 \leq p \leq n \\ F^p \oplus \bar{F}^{n-p+1} \simeq V_{\mathbb{C}} \end{array} \right.$$

$$\left\{ \begin{array}{l} V^{p,q} = F^p \oplus \bar{F}^q = \bar{V}^{q,p} \\ V_{\mathbb{C}} = \bigoplus V^{p,q} \\ F^p = \bigoplus_{p' \geq p} V^{p',q} \end{array} \right.$$

Ex $H^n(X, \mathbb{Q})$

Polarized Hodge structure of weight n

(V, Q, F^p)

$$\left\{ \begin{array}{l} Q(F^p, F^{n-p+1}) = 0 \end{array} \right. \quad (I)$$

$$\left\{ \begin{array}{l} i^{p-q} Q(V^{p,q}, \bar{V}^{p,q}) > 0 \end{array} \right. \quad (II)$$

Ex $H^n(X, \mathbb{Q})_{\text{prim}}$

Mixed Hodge structure

$$(V, W, F^p)$$

$$\left\{ \begin{array}{l} W = \text{increasing filtration} / \mathbb{Q} \\ G_m^W V = W_m V / W_{m-1} V \\ F^p \text{ induces on } G_m^W V \text{ a HS of weight } m \end{array} \right.$$

Ex Σ complete, $V = H^m(\Sigma, \mathbb{Q})$
and $0 \leq m \leq n$

Ex Σ smooth and affine, $V = H^m(\Sigma, \mathbb{Q})$
and $n \leq m \leq 2n$

Period domain

$$D = \{ F' : (V, Q, F') = \text{PHS} \}$$

$$D = G_{\mathbb{R}} / H, \quad G_{\mathbb{R}} = G(\mathbb{R}) \text{ where} \\ G = \text{Aut}(V, Q) \text{ and} \\ H = \text{isotropy of } F_0 \in D$$

Ex $n=1$ $V_{\mathbb{C}} = V^{1,0} \oplus \bar{V}^{1,0}$
 $Q(V^{1,0}, V^{1,0}) = 0, \perp Q(V^{1,0}, \bar{V}^{1,0}) > 0$

-3-

$$u \rightarrow D = S_V(2g, \mathbb{R}) / U(g) = \mathcal{H}_g$$

$$\underline{\text{Ex: } n=2} \quad V_G = \underbrace{V^{2,0}}_{\dim a} \oplus \underbrace{V^{1,1}}_{\dim b} \oplus \overline{V}^{2,0}$$

$$u \rightarrow D \cong SO(2a, b) / U(a) \times SO(b)$$

Compact dual

$$\tilde{D} = \{ F : Q(F^p, F^{n-p+1}) = 0 \}$$

$$\tilde{D} = G_G / \mathcal{P}, \quad \mathcal{P} = \text{parabolic}$$

$$u \rightarrow D = \text{open } G_{\mathbb{R}}\text{-orbit in } \tilde{D}$$

$G_{\mathbb{R}}$ -orbit structure of $\tilde{D}$ is a rich subject

Period mapping

$$\cdot \quad X \xrightarrow{\pi} B, \quad \pi^{-1}(b) = \Sigma_b \text{ smooth}$$

• monodromy representations


$$\varrho: \pi_1(B) \rightarrow \text{Aut}(H^n(\Sigma_{b_0}, \mathbb{Q})_{\text{prim}}, \mathbb{Q})$$

$$u \rightarrow \underline{\Gamma}: B \rightarrow \Gamma \backslash D$$

$$\text{where } \varrho(\pi_1(B)) < \Gamma < G$$

Singularities (I)

$$\begin{array}{ccc} \mathcal{X} & \subset & \overline{\mathcal{X}} \\ \downarrow & & \downarrow \\ \Delta^* & \subset & \Delta \end{array}$$



$$T = T_S T_u$$

↑
semi-simpl

↑
irreducible

$$T^m = I \quad (\text{eigenvalues of } T \text{ are roots of unity})$$

$$T_u = e^N, \quad N \text{ nilpotent and } N^{n+1} = 0$$

Limiting mixed Hodge structure (LMHS)

$$\cdot \text{ nilpotent } N \text{ gives unique } W(N)$$

$$\cdot N: W_m(N) \rightarrow W_{m-2}(N)$$

$$\cdot N^k: Gr_{n+k}^{W(N)} V \cong Gr_{n-k}^{W(N)} V$$

$$\cdot \text{ there exists } \mathfrak{sl}_2 = \{N, H, N^+\} \text{ such that } N = \partial_x$$

$$LMHS = \{V, W(N), F_{\text{lim}}\}$$

$$\text{Ex: } \mathbb{P}: \Delta^* \rightarrow \{T^{\mathbb{Z}}\} \setminus \{D\} \text{ gives LMHS (Schmid)}$$

Local monodromy cycle theorem

$$f_t: X_t \rightarrow X_0$$



$$\mapsto f^* H^*(X_0) = \text{loc} N$$

Several variable case

$$\Delta^{\frac{1}{2}} \times \Delta^l \rightarrow \Gamma_{bc} \setminus D$$

- $W(N_\lambda)$ independent of λ when

$$N_\lambda = \sum \lambda_i N_i, \quad \lambda_i > 0$$

- relative weight filtration property

- asymptotics of degenerating PHS's
(Cattani - Kaplan - Schrieber)